

Research

Open Access

6-sulfo sialyl Lewis X is the common receptor determinant recognized by H5, H6, H7 and H9 influenza viruses of terrestrial poultry

Alexandra S Gambaryan¹, Alexander B Tuzikov², Galina V Pazynina², Julia A Desheva³, Nicolai V Bovin², Mikhail N Matrosovich^{*4} and Alexander I Klimov⁵

Address: ¹Chumakov Institute of Poliomyelitis and Viral Encephalitis, RAMS, 142782 Moscow, Russia, ²Shemyakin Institute of Bio-organic Chemistry, RAS, 117997 Moscow, Russia, ³Institute of Experimental Medicine, RAMS, 197376 St. Petersburg, Russia, ⁴Institute of Virology, Philipps University, 35043 Marburg, Germany and ⁵Influenza Division, Centers for Disease Control and Prevention, Atlanta, GA 30333, USA

Email: Alexandra S Gambaryan - alexandra.gambaryan@gmail.com; Alexander B Tuzikov - tuzikov@carb.ibch.ru; Galina V Pazynina - pazynina@carb.ibch.ru; Julia A Desheva - desheva@ok.ru; Nicolai V Bovin - bovin@carb.ibch.ru; Mikhail N Matrosovich^{*} - M.Matrosovich@gmail.com; Alexander I Klimov - aklimov@cdc.gov

^{*} Corresponding author

Published: 24 July 2008

Received: 3 May 2008

Virology Journal 2008, 5:85 doi:10.1186/1743-422X-5-85

Accepted: 24 July 2008

This article is available from: <http://www.virologyj.com/content/5/1/85>

© 2008 Gambaryan et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Influenza A viruses of domestic birds originate from the natural reservoir in aquatic birds as a result of interspecies transmission and adaptation to new host species. We previously noticed that influenza viruses isolated from distinct orders of aquatic and terrestrial birds may differ in their fine receptor-binding specificity by recognizing the structure of the inner parts of Neu5Ac α 2-3Gal-terminated sialyloligosaccharide receptors. To further characterize these differences, we studied receptor-binding properties of a large panel of influenza A viruses from wild aquatic birds, poultry, pigs and horses.

Results: Using a competitive solid-phase binding assay, we determined viral binding to polymeric conjugates of sialyloligosaccharides differing by the type of Neu5Ac α -Gal linkage and by the structure of the more distant parts of the oligosaccharide chain. Influenza viruses isolated from terrestrial poultry differed from duck viruses by an enhanced binding to sulfated and/or fucosylated Neu5Ac α 2-3Gal-containing sialyloligosaccharides. Most of the poultry viruses tested shared a high binding affinity for the 6-sulfo sialyl Lewis X (Su-SLe^x). Efficient binding of poultry viruses to Su-SLe^x was often accompanied by their ability to bind to Neu5Ac α 2-6Gal-terminated (human-type) receptors. Such a dual receptor-binding specificity was demonstrated for the North American and Eurasian H7 viruses, H9N2 Eurasian poultry viruses, and H1, H3 and H9 avian-like virus isolates from pigs.

Conclusion: Influenza viruses of terrestrial poultry differ from ancestral duck viruses by enhanced binding to sulfated and/or fucosylated Neu5Ac α 2-3Gal-terminated receptors and, occasionally, by the ability to bind to Neu5Ac α 2-6Gal-terminated (human-type) receptors. These findings suggest that the adaptation to receptors in poultry can enhance the potential of an avian virus for avian-to-human transmission and pandemic spread.

Background

The recent pandemic threat caused by the widespread circulation of H5N1 avian influenza viruses and their occasional transmission to humans as well as human infections caused by chicken H9N2, H7N7 and H7N3 viruses highlighted the need for a detailed study of host restriction mechanisms of influenza viruses. Numerous studies support the concept that alteration of the receptor specificity of an avian virus is essential for its transmission into humans as well as for human-to-human transmission and pandemic spread (reviewed in ref. [1,2]).

The history of research into the receptor binding phenotypes of influenza viruses can be divided into two periods: before and after 1997 when first human infections with chicken H5N1 viruses were documented. Before 1997, it was established that human influenza viruses recognize Neu5Ac α 2-6Gal-terminated receptors, avian viruses recognize Neu5Ac α 2-3Gal-terminated receptors while swine viruses recognize both of them [3-8]. It was shown that the receptor-binding site (RBS) of the hemagglutinin (HA) of avian viruses is evolutionally very stable. In addition to eight amino acids forming the HA RBS, which are conserved in all influenza A viruses (positions 97, 98, 134, 139, 153, 183, 184 and 195; H3 numbering is used here and throughout the paper), there are six more amino acids conserved in HAs of duck viruses (positions 138, 190, 194, 225, 226 and 228), and these are positions where human HAs are different from duck viruses [7]. Virus receptor binding specificity was found to correlate with the level of expression of relevant sialic acids determinants on the target cells of different host species. Thus, epithelial cells of human airway epithelium were shown to express high amounts of Neu5Ac α 2-6Gal-terminated sialyloligosaccharides, duck intestinal epithelium predominantly contains Neu5Ac α 2-3Gal-terminated receptors while swine tracheal epithelium contains both receptor types [8,9]. It was hypothesized that alteration of receptor specificity of avian viruses in some intermediate host, such as swine, might facilitate their transmission to humans [10].

After 1997, it became clear that avian H5N1 viruses are capable of replicating in humans [11,12] despite their avian-virus-like preference for Neu5Ac α 2-3Gal-containing receptors and lack of binding to human-type receptors [13]. It was shown afterwards that human airway epithelial cells express 2-3-linked sialic acid receptors with a density sufficient for the entry and replication of avian viruses [14,15].

Furthermore, a Eurasian lineage of poultry H9N2 viruses was discovered, which recognized Neu5Ac α 2-6Gal-terminated sialyloligosaccharides, thus indicating that some avian influenza viruses may display a human-virus-like

receptor specificity [16-18]. It was also demonstrated that chicken and quail intestinal cells contain both Neu5Ac α 2-3Gal and Neu5Ac α 2-6Gal sialyloligosaccharides, in contrast to duck cells that contain only Neu5Ac α 2-3Gal [19-23].

Although the Neu5Ac α 2-3Gal receptor specificity is shared by the majority of avian viruses, viruses adapted to different avian species can differ in their ability to recognize the third saccharide and more distant moieties of Neu5Ac α 2-3Gal-terminated receptors. For example, duck viruses of various subtypes preferentially bound to glycoprotein O-chain trisaccharide Neu5Ac α 2-3Gal β 1-3GalNAc α , whereas H5N1 chicken viruses preferred receptors with inner β -N-acetylglucosamine moiety, Neu5Ac α 2-3Gal β 1-4GlcNAc β [20]. Sulfation of the saccharide core produced no effect on binding of duck viruses, whereas chicken and human viruses isolated in 1997 in Hong Kong demonstrated an extraordinarily high affinity for sulfated trisaccharide Neu5Ac α 2-3Gal β 1-4(6-HSO₃)GlcNAc (Su-3'SLN) [24,25].

In the present study, we characterized the receptor-binding specificity of a broad set of influenza A viruses from wild aquatic birds, poultry, pigs and horses.

Results

Receptor-binding specificity

To determine the receptor-binding specificity of avian and mammalian influenza viruses, we tested their binding to 9 distinct polymeric glycoconjugates (see Fig. 1 and Table 1 for structural formulas and abbreviations). One of the glycoconjugates harboured 6-linked sialyloligosaccharide, Neu5Ac α 2-6Gal β 1-4GlcNAc (6'SLN). The oligosaccharide parts of the other glycopolymers shared the same terminal Neu5Ac α 2-3Gal moiety but differed: (i) by the type of the bond between galactose and the next sugar residue (β 1-3 or β 1-4), (ii) by the nature of this residue (GlcNAc β or GalNAc α), and (iii) by constituents at different positions on the GlcNAc ring (fucose or/and sulfo group). All studied oligosaccharide structures have been found in natural glycoproteins or glycolipids [26]. Virus binding to glycoconjugates was determined in a competitive solid-phase assay and expressed in terms of binding affinity constants (Fig. 2).

Each of the tested viruses bound SLec, Su-SLec and STF with the same affinity and none of the viruses discriminated between SLex and SLea. We, therefore, do not show here the binding data for Su-SLec, STF and SLea. The patterns of viral binding to the panel of receptor analogues varied significantly among viruses of different subtypes and host species (Fig. 2), however, several distinctive groups of viruses with typical receptor binding phenotypes could be recognized as described below.




Figure 1
Molecular models of sialyloligosaccharides. The models depict sialyloligosaccharide parts of glycopolymers that were tested for their binding to influenza viruses. Corresponding structural formulas are given in the Table 1. The figures were generated using Discovery Studio ViewerPro5.0 software (Accelrys Inc.).

Viruses of various subtypes isolated from wild ducks
 These viruses displayed the highest binding affinity for glycoconjugates with $\beta(1-3)$ linkage between Neu5Ac α 2-3Gal disaccharide fragment and the next GlcNAc residue, i.e., SLe^c, Su-SLe^c and STF. Other characteristic features of duck viruses were their low affinity for fucosylated sialyloligosaccharides SLe^a and SLe^x, nearly equal affinity for sulfated and non-sulfated sialyloligosaccharides, and a lack of appreciable binding to 6'SLN.

Sialyloligosaccharide	Abbreviation
Neu5Ac α 2-3Gal β 1-4GlcNAc β	3'SLN
Neu5Ac α 2-3Gal β 1-4(6-HSO ₃)GlcNAc β	Su-3'SLN
Neu5Ac α 2-3Gal β 1-4(Fuc α 1-3)GlcNAc β	SLe ^x
Neu5Ac α 2-3Gal β 1-4(Fuc α 1-3)-(6-HSO ₃)GlcNAc β	Su-SLe ^x
Neu5Ac α 2-3Gal β 1-3GlcNAc β	SLe ^c
Neu5Ac α 2-3Gal β 1-3(6-HSO ₃)GlcNAc β	Su-SLe ^c
Neu5Ac α 2-3Gal β 1-3GalNAc α	STF
Neu5Ac α 2-3Gal β 1-3(Fuc α 1-4)GlcNAc β	SLe ^a
Neu5Ac α 2-6Gal β 1-4GlcNAc	6'SLN

Viruses with H6 HA
 Five viruses with H6 HA tested in this study were isolated from different avian species (turkey, shearwater, teal, chicken and gull). Unlike typical duck viruses, all H6 viral isolates efficiently bound to fucosylated sialyloligosaccharides SLe^x and SLe^a.

Viruses with H7 HA
 Viruses of H7 subtype from two evolutionary lineages were tested: 1) American avian H7N2 viruses and closely related human isolate A/New York/107/03 (H7N2) [27], and 2) Eurasian H7N7 human isolates that were transmitted to humans from infected poultry during the 2003 outbreak in the Netherlands [28]. Viruses from both lineages showed enhanced binding to sulfated sialyloligosaccharides with the Gal β 1-4GlcNAc β core. The American viruses displayed the highest affinity for Su-3'SLN, whereas the H7N7 viruses from the Netherlands had particularly high affinity for Su-SLe^x. It was found unexpectedly that all H7 viruses tested displayed moderate binding affinity for human-type receptor 6'SLN (Fig. 2).

H9N2 viruses
 The H9N2 viruses tested could be arbitrarily separated into three groups, North American viruses and distantly

Virus		Sialylglycoconjugate					
		3'SLN	Su-3'SLN	SLe ^x	Su-SLe ^x	SLe ^c	6'SLN
Duck viruses							
Duck /Hong Kong/278/78	H2N9	20	10	>50	>50	10	>5000
Duck/Nanchang/2-0485/00	H2N9	20	10	>50	>50	10	>5000
Duck/Buryatia/652/88*	H3N8	10	10	50	50	5	>5000
Mallard/New York/670/78	H4N6	4	8	100	>200	2	>5000
Duck /Buryatiya/1905/00*	H4N6	20	20	100	100	10	>5000
Duck/Primorie/3628/02*	H9N2	10	8	100	30	6	>5000
Mallard/Netherlands/02/00	H10N4	20	7	>50	30	5	>5000
H6							
Turkey/Massachusetts/65	H6N2	20	30	20	30	10	>5000
Shearwater/Australia/1/72	H6N5	20	30	10	30	15	>5000
Teal/Hong Kong/W312/97	H6N1	50	>50	20	>50	20	>5000
Chicken/New York/13237/98	H6N8	7	10	7	20	5	>5000
Gull/Moscow/3100/06	H6N2	5	5	2	2	3	>5000
H7							
Turkey/Virginia/4529/02	H7N2	2	1	20	4	10	200
Avian/New York/273874/03	H7N2	10	2	30	8	20	500
Chicken /NJ/294598-12/04	H7N2	5	1	20	5	5	200
Chicken /Delaware/296763/04	H7N2	15	5	30	15	20	200
New York/107/03	H7N2	10	2	>100	15	20	200
Netherlands/219/03	H7N7	4	1	5	0.5	5	500
Netherlands/230/03	H7N7	4	1	5	0.5	5	500
Netherlands/231/03	H7N7	4	1	5	0.5	5	500
H9N2							
Goose/Minnesota/5773/80	H9N2	20	10	50	20	10	>5000
Turkey/Wisconsin/1/66	H9N2	5	5	20	30	2	>5000
Turkey/Minnesota/38391-6/95	H9N2	10	10	10	20	5	>5000
Pheasant/Wisconsin/1780/88	H9N2	10	3	2	1	10	>5000
Chicken/New Jersey/12220/97	H9N2	5	5	0.5	0.5	5	>5000
Chicken/Korea/96323/96	H9N2	10	3	30	10	10	>5000
Hong Kong/1073/99	H9N2	>100	>100	>100	>100	>100	20
Quail/Hong Kong/G1/97	H9N2	>100	>100	20	20	>100	20
Chicken/Hong Kong/FY20/99	H9N2	>100	3	>100	5	>100	50
Duck/Hong Kong/Y280/97	H9N2	>100	0.3	>100	0.1	>100	10
Chicken/Hong Kong/G9/97	H9N2	>100	5	>100	10	>100	50
Chicken/Hong Kong /SF3/99	H9N2	>100	3	>100	5	>100	50
Hong Kong/2108/03	H9N2	>100	10	>100	10	>100	50
Swine							
Swine/Hong Kong/9/98	H9N2	>100	1	>100	2	>100	50
Swine/Finistere/2899/82*	H1N1	30	30	20	5	30	100
Swine/France/80*	H1N1	30	20	40	10	40	100
Swine/Kazakhstan/48/82*	H3N6	10	3	40	3	15	1000
Equine							
Equine/Kentucky/5/02	H3N8	3	1	10	1	4	>5000
Equine/Ohio/1/03	H3N8	3	0.5	5	0.4	4	>5000
Canine/Florida/43/04	H3N8	3	0.5	5	0.4	4	>5000
H5N1							
Chicken/ Hong Kong /220/97	H5N1	3	0.5	80	20	10	>5000
Chicken/Vietnam/NCVD11/03	H5N1	1	0.3	20	1	2	>5000
Human pandemic viruses							
USSR/039/68	H3N2	>200	>200	>200	>200	>200	4
Canada/228/68	H3N2	>200	>200	>200	>200	>200	2

Figure 2

Binding affinity constants of virus complexes with sialylglycopolymers. The constants were determined as described in the Methods and were expressed in μM of sialic acid. Higher values of constants correspond to lower binding affinities. The data were averaged from 3 sets of experiments. Standard errors did not exceed 50% of the mean values. Viruses labelled with asterisk were kindly provided by Dr. S. Yamnikova, the Ivanovsky Institute of Virology, Moscow, Russia. Colours depict relative levels of binding for each individual virus: red – maximal binding; yellow – good binding; pale cyan – weak binding; blue – no detectable binding.

related virus A/Chicken/Korea/96323/96 [29] and two evolutionary lineages of poultry viruses from Southeast Asia, G1 and G9 [16-18].

Receptor-binding affinity of A/Goose/Minnesota/5773/80 and A/Turkey/Wisconsin/1/66 was similar to that of A/Duck/Primorie/3628/02 (H9N2) and duck viruses of other subtypes: they preferentially bound to SLe^c and bound poorly to fucosylated sialyloligosaccharides. Other North American viruses such as A/Chicken/New Jersey/12220/97 and A/Pheasant/Wisconsin/1780/88 showed high binding affinity for SLe^a, SLe^x and Su-SLe^x. A/Chicken/Korea/96323/96 had an increased affinity for sulfated sialyloligosaccharides Su-3'SLN and Su-SLe^x. None of these viruses bound 6'SLN.

In contrast to North American viruses, Asian isolates from G1- and G9- lineages bound to 6'SLN. The binding pattern of the human isolate A/Hong Kong/1073/99 (H9N2) resembled that of pandemic human viruses A/USSR/039/68 and A/Canada/228/68 (see Fig. 2, bottom lines): all these three viruses strongly bound to 6'SLN and did not appreciably bind to Neu5Acα2-3Gal-containing oligosaccharides. A/Quail/Hong Kong/G1/97 demonstrated high affinity for 6'SLN, SLe^x and Su-SLe^x and did not bind to any of non-fucosylated Neu5Acα2-3Gal-containing receptors. All other Asian poultry viruses tested displayed moderate binding to 6'SLN, bound much stronger to sulfated receptors Su-3'SLN and Su-SLe^x and did not bind at all to 3'SLN, SLe^c, SLe^x, and Su-SLe^c.

Swine viruses

Four viruses isolated from pigs were tested. A/Swine/Hong Kong/9/98 belonged to the G9 clade of H9N2 viruses, A/Swine/Finistere/2899/82 and A/Swine/France/80 represented the European avian-like swine virus lineage and A/Swine/Kazakhstan/48/82 was a sporadic avian-like H3N6 isolate. A common feature of these viruses was their high affinity for Su-SLe^x and a moderate affinity for 6'SLN.

Equine viruses

Equine H3N8 viruses including the equine-like canine isolate A/Canine/Florida/43/2004 [30] showed a strong binding affinity for Neu5Acα2-3Gal receptors, preferring sulfated ones, Su-3'SLN or Su-SLe^x.

H5N1 Asian viruses

Viruses of this group were extensively analyzed in our previous studies [24,25]. Two typical chicken isolates were tested here for a comparison with other poultry viruses. Both A/Chicken/Hong Kong/220/97 and A/Chicken/Vietnam/NCVD11/03 revealed increased affinity for Su-3'SLN. The latter virus in addition showed a high affinity for Su-SLe^x.

Analysis of HA amino acid sequences and molecular modelling of the complexes of Su-SLe^x with H3, H7 and H9 HA

The characteristic feature of the duck viruses tested herein was their poor binding to fucosylated sialyloligosaccharides (Fig. 2, upper part). This receptor-binding phenotype agreed with that described earlier for a variety of viruses from wild ducks [20,24,31]. In order to understand the molecular basis of this phenotype, we modelled a putative disposition of the fucosylated receptor Su-SLe^x in the receptor-binding site of the HA of Duck/Ukraine/1/63 (H3N8) [32]. The modelling predicted that the fucose moiety would come into a significant sterical conflict with the side chain of Trp222 (Fig 3). We next compared the HA sequences of more than 400 duck influenza viruses of H1, H2, H3, H4, H5, H8, H9, H10, H11 and H14 subtypes available from the Genbank. All of these viruses had a bulky amino acids (Arg, Lys, Trp, Leu, or Gln) in position 222 of the HA. We suggest on this basis that partial overlap of the fucose moiety with the bulky amino acid in position 222 could be a universal mechanism that reduces the capability of duck viruses to bind fucosylated receptors.

Our analysis of 68 published H6 HA sequences revealed that 67 of them have Ala222. This finding suggests that a relatively good binding of H6 viruses to fucosylated sialyloligosaccharides SLe^x and SLe^a (Fig. 2) could be explained by a lack of interference between the fucose moiety and the short side chain of the alanine in position 222 of the HA. Essential role of amino acid in position 222 in the binding of fucosylated receptors was also supported by the comparison of HA sequences of the H9N2 viruses, A/Goose/Minnesota/5773/80 and A/Chicken/New Jersey/12220/97 (Fig. 2 and Fig. 4). The latter virus had His222 and bound SLe^x 100-times better than the former virus (Leu222).

The high binding affinity of H7 viruses to sulfated sialyloligosaccharides suggested that the sulfo group interacts with some charged amino acid residue in the receptor-binding site. To test this possibility, we modelled potential contacts of Su-SLe^x with the receptor-binding pocket of avian H3 and H7 HAs [32,33]. In the case of H3 duck virus, the sulfo group faced towards solution and did not form obvious direct contacts with the protein. However, in the case of the H7 HA, the sulfo group of Su-SLe^x was located in a close proximity to the side chain of Lys193, which is highly conserved among viruses with H7 HA (Fig 3). This finding suggests that enhanced affinity of H7 viruses for Su-3'SLN and Su-SLe^x is due to favourable charged interactions between the sulfo group of the receptor and amino group of Lys193. The same mechanism is likely responsible for the high affinity for Su-3'SLN of H5N1 (Gambaryan et al., 2004, 2006) and H3N8 equine




Figure 3
Models of complexes of H3, H7 and H9 hemagglutinins with Su-SLe^x. The models were generated as described in the Methods using the crystal structures of the HAs of the viruses A/Duck/Ukraine/1/63 (H3N8) [32], A/Turkey/Italy/02 (H7N1) [33] and A/Swine/Hong Kong/9/98 (H9N2) [37]. The fucose moiety and the sulfo group of Su-SLe^x are shown as mesh surfaces. Amino acid residues described in the text are numbered.

viruses (Fig. 2) since viruses of both these groups have lysine in position 193.

South-eastern Asian H9N2 viruses have multiple amino acid substitutions in the receptor-binding region of the HA, most notably, the mutation Gln226Leu [34] (Fig. 4). The typical human-virus-like receptor specificity of A/Hong Kong/1073/99 (H9N2) is in a good agreement with

the notion that a single Gln226Leu replacement shifts the receptor specificity from recognition of Neu5Ac α 2-3Gal to recognition of Neu5Ac α 2-6Gal receptors [35,36]. A/Quail/Hong Kong/G1/97 (H9N2) differs from A/Hong Kong/1073/99 (H9N2) by the Gly225Asp substitution (Fig. 4) that markedly enhances the affinity of this virus for fucosylated 3-linked receptors SLe^x and Su-SLe^x and leads to a rather an unusual receptor-binding phenotype (Fig. 2).

The H9N2 viruses of G9-lineage harbour substitutions Gln226Leu and Glu190Ala/Thr/Val in the HA [18,34] (Fig. 4). Viruses with Ala or Thr in position 190 bound Su-3'SLN and Su-SLe^x with the highest affinity and demonstrated moderate affinity for 6'SLN (Fig. 2). As these viruses did not noticeably bind to Su-SLe^c, specific orientation of the sulfo group rather than its negative charge alone seems to be essential for the binding. We used the crystal structure of the HA of A/Swine/Hong Kong/9/98 (H9N2) (G9-lineage) in complex with 3-linked receptor [37] for the modelling of H9 HA interactions with Su-SLe^x (Fig. 3). Due to the amino acid substitutions in positions 226 and 190 of the H9 HA, the conformation of 3-linked galactose in this complex differs from that in the H3 avian HA [32,37], leading to corresponding differences in the putative disposition of Su-SLe^x (compare H3 and H9 complexes in Fig. 3). In the H9 HA, the fucose moiety shifts upwards resolving the steric interference with amino acid in position 222, whereas the sulfo group shifts downwards and fits into a cavity formed by amino acids in positions 190 and 186 and by the solvent water molecules bound to residues 98, 228 and 227 (PDB:1JSD[37]). This could explain why the substitutions Gln226Leu and Glu190Val in the H9 HA, that increased virus affinity for Neu5Ac α 2-6Gal, at the same time significantly enhanced its affinity for sulfated Neu5Ac α 2-3Gal-containing receptors, Su-3'SLN and Su-SLe^x.

Discussion

Although almost all avian viruses use the same terminal disaccharide Neu5Ac α 2-3Gal as receptor, the evolution of distinct virus lineages adapted to distinct avian species (wild ducks, gulls, or terrestrial poultry) has led to specialized abilities to recognize longer oligosaccharide chains. Thus, duck viruses have the highest affinity for SLe^c and STF (Neu5Ac α 2-3Gal β 1-3GalNAc α). We demonstrated earlier that duck viruses bind strongly to gangliosides from duck intestine as well as to GD1a ganglioside, which is terminated by Neu5Ac α 2-3Gal β 1-3GalNAc β [7,19]. It is possible that gangliosides with this termination serve as functional receptors of influenza viruses in the duck intestine.

Our present study indicated that receptor specificity of viruses from different lineages adapted to quail and chicken differed from that of wild duck viruses. Sulfated

	*	180	*	200	*	220	*
Duck/Alberta/321/88		QDAQYTNNEGKNILFMWGIHHPPTDTE		QTNLYKKADTTTSVTTEDINRTFKPVIGPRPLVNGQQGRIDYY			
Duck/Primorie/3628/02		
Goose/Minnesota/5773/80		
Turkey/Wisconsin/1/66		.N.H.....RE.....D...N.....		
Turkey/Minnesota/38391/95	T...V.....ND...T.....M...P.....		
Chicken/N.Jersey/12220/97	R.....			H.....	
Chicken/Korea/96323/96	DW.....		I.....I.....		
Hong Kong/1073/99	R.S...V.....Y.....		IRN.....L.....		L.....	
Quail/Hong Kong/G1/97	R.S...V.....Y.....		IRN.....L.....		DL.....	
Chicken/Hong Kong/G9/97	R.S.....N.....A.....		TRT.....A.....		L.....	
Chicken/Hong Kong/FY20/99	R.....N.....A.....		TRT.....A.....		L.....N.....	
Duck/Hong Kong/Y280/97	R.....N.....T.....		TRT.....A.....		L.....	
Hong Kong/2108/03	R.S.....N.....V.....		TRT.....A.....		L.....	
Swine/Hong Kong/9/98	R.S.....N.....V.....		TRT.....		LH.....	

Figure 4

Partial HA amino acid sequences of H9N2 viruses. The sequences were obtained from GenBank. Differences with respect to the top sequence are shown. Amino acids in positions 190, 222, and 226 are highlighted. The figure was generated with GeneDoc 2.6 software [50].

and fucosylated 3'SLN is a suitable receptor for most of these poultry viruses. It was shown recently that bi-antennary a2-6/3 sialylated glycans with Gal β 1-4GlcNAc β core are major sialylated *N*-glycans expressed by intestinal epithelial tissues in both chicken and quail [23]. This fact is in accord with preferential binding of quail and chicken viruses to sialyloligosaccharides with Gal β 1-4GlcNAc β core.

Gull viruses appear to be adapted to fucosylated receptors, such as SLe^x [31]. We suggested earlier that the presence of glycine in HA position 222 of H13 and H16 viruses is essential for this binding phenotype [38]. In this study, we found that all tested H6 viruses, similarly to gull viruses, demonstrated enhanced affinity for SLe^a and SLe^x and that alanine in position 222 of the H6 HA is likely to play an essential role in this specificity. Since almost all sequenced H6 HA have Ala222, viruses of this subtype should be able to recognize receptor determinants that are optimal for duck (SLe^c), gull (SLe^x) and chicken (Su-SLe^x) viruses. This feature of H6 viruses would agree with their known promiscuous host range [39]. Some American H9 poultry viruses with substitution in position 222 showed receptor binding phenotype that was similar to that of H13, H16 and H6 viruses. It could be speculated that mutations in position 222 that improve sterical accommodation of the fucose moiety could represent one general pathway for adaptation of duck viruses to fucosylated receptors present in gulls and chickens.

Asian H5N1 viruses, H7 poultry viruses and equine H3N8 viruses realized another pathway of adaptation to recognition of the Su-SLe^x determinant via the favourable electrostatic interactions between the sulfo group and the amino group of Lys193.

One more pathway of viral adaptation to Su-SLe^x can be achieved through a substitution of conserved glutamic acid in the HA position 190. Importantly, this substitution that leads to enhanced binding to Su-SLe^x is often accompanied by the enhanced viral binding to the human-type receptor 6'SLN. Thus, high affinity for Su-SLe^x and moderate affinity for 6'SLN was detected in this study for G9-like H9N2 viruses, and previously for H1N1 swine [40] and human [41] viruses. H7 viruses with high affinity for Su-SLe^x also showed detectable binding to 6'SLN (Fig. 2).

It is not clear whether the ability of H7 and H9 poultry viruses to bind to 6'SLN provides them with some evolutionary advantage. For example, the binding of these viruses to 6'SLN does correlate with the presence of 6'SLN-containing receptors in epithelial tissues of gallinaceous birds [19-23]. Alternatively, the ability of poultry viruses to bind to 6'SLN could be an accidental consequence of their adaptation for the binding to Su-SLe^x due to some sterical similarity between Su-SLe^x and 6'SLN in the regions of Neu5Ac-Gal glycosidic linkage and of the NAc-moiety of the GlcNAc residue (Fig. 1).

The binding data (Fig. 2) show that the receptor specificity of poultry H5, H7, and H9 viruses is similar to that of equine and avian-like swine viruses. If sulfated and fucosylated sialyloligosaccharides are present in the target cells of both terrestrial poultry and mammals, the adaptation of aquatic bird viruses to poultry could facilitate their replication in mammals, including humans.

Conclusion

It is generally believed that alteration of the receptor specificity is a prerequisite for the highly effective replication and human-to-human transmission which characterize

pandemic influenza viruses [1,2,42]. Here we found that several independent lineages of poultry influenza viruses differ from their precursors in aquatic birds by enhanced binding to 6-sulfo sialyl Lewis X and that this binding specificity is accompanied by the ability of the virus to bind to human-type receptor 6'SLN. We therefore suggest that the adaptation to Su-SLe^x receptor in terrestrial poultry could enhance the potential of an avian virus for avian-to-human transmission and pandemic spread.

Methods

Materials

Oligosaccharides conjugated with polyacrylamide (~30 kDa) were synthesized from spacers sialyloligosaccharides (spacer = -OCH₂CH₂CH₂NH₂ or -NHCOCH₂NH₂) and poly(4-nitrophenylacrylate) having m.w. 30 kDa by the method described earlier [43,44]. Spacers oligosaccharides were synthesized as described previously [45-47].

Viruses

The majority of viruses in the study were from the repository of the Influenza Division, CDC, USA; some isolates were from the collection of the D.I. Ivanovsky Institute of Virology, Moscow. Viruses were grown in 9-day-old embryonated chicken eggs and were inactivated by treatment with beta-propiolactone as described previously [13]. The allantoic fluids were clarified by low-speed centrifugation; the viruses were pelleted by high-speed centrifugation, resuspended in 0.1 M NaCl, 0.02 M Tris buffer (pH 7.2) containing 50% glycerol, and stored at -20°C.

The binding affinity of influenza viruses for sialylglycoconjugates

Receptor specificity of influenza viruses was evaluated in a competitive assay based on the inhibition of binding to solid-phase immobilized virus with bovine fetuin labelled with horseradish peroxidase [48]. The competitive reaction was performed at 2-4 °C for 30 min in PBS with 0.01% of Tween-20; 0.05% of BSA and 3 µM of the sialidase inhibitor 4-amino-Neu5Ac-en. The data were expressed in terms of affinity constants (K_{aff}) formally equivalent to the dissociation constants of virus-receptor complexes. For the calculation of the constants, concentration of the sialic acid residues in the solution was used. Each set of experiments presented in the Fig. 2 was repeated three-four times with similar results. Data were averaged from 3 sets of experiments.

Molecular models

Atomic coordinates of SLe^x (PDB:2KMB) [49], H7 HA (PDB:1TI8) [33], H9 HA (PDB:1JSD) [37] and H3 and H9 HA complexes with NeuAc α 2-3Gal-containing pentasaccharide LSTa (PDB:1MQM and PDB:1JSH) [32,37] were obtained from Brookhaven Protein Data Bank. The

molecular models were generated using DS ViewerPro 5.0 software (Accelrys Inc.).

The model of Su-SLe^x was constructed on the basis of SLe^x structure (PDB:2KMB), by replacing the hydrogen atom of the 6-OH group of GlcNAc by HSO₃ group.

The models of Su-SLe^x in the receptor-binding sites of H3 and H9 HA were made by superimposing the galactose residue of the Su-SLe^x over the galactose residue of LSTa. The model of Su-SLe^x in the receptor-binding site of H7 HA was generated by superimposing the protein chain of the H3 HA complex with Su-SLe^x over the protein chain of H7 HA (PDB:1TI8). The OH groups of Tyr98, SG atoms of Cys139, CZ3 atoms of Trp153, CD atoms of Glu190 and CA atoms of Tyr 195 were used to align two proteins.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Conception and design of the study, manuscript preparation (ASG, NVB, AIK, MNM); experimental work (ASG, ABT, GVP, JAD); co-ordination of the study (AIK, NVB).

Disclaimer

The findings and conclusions in the report are those of the authors and do not necessarily represent the views of the funding agencies.

Acknowledgements

We thank Dr. Svetlana Yamnikova (D.I. Ivanovsky Institute of Virology, Moscow, Russia) for providing us with duck influenza viruses and Dr. Michael Shaw (Influenza Division, Centers for Disease Control and Prevention, Atlanta, GA, USA) for critical review of the paper. This study was supported by research grants 05-04-48934 from the Russian Foundation for Basic Research, RAS Presidium program 'Molecular and Cell Biology', ISTC grant No:5 2464 and the European Commission projects FLUPATH, FLUINNATE and FLUVACC.

References

1. Matrosovich MN, Klenk HD, Kawaoka Y: **Receptor specificity, host range and pathogenicity of influenza viruses.** In *Influenza Virology: Current Topics* Edited by: Kawaoka Y. Wymondham: Caister Academic Press; 2006:95-137.
2. Klenk HD, Matrosovich M, Stech J: **Avian Influenza – Molecular Mechanisms of Pathogenesis and Host Range.** In *Molecular Biology of Animal Viruses* Edited by: Mettenleiter T, Sabirina F. United Kingdom: Caister Academic Press; 2008:253-301.
3. Paulson JC: **Interaction of animal viruses with cell surface receptors.** In *The Receptors Volume 2*. Edited by: Corn M. Orlando: Academic Press; 1985:131-219.
4. Rogers GN, D'Souza BL: **Receptor binding properties of human and animal H1 influenza virus isolates.** *Virology* 1989, **173**:317-322.
5. Connor RJ, Kawaoka Y, Webster RG, Paulson JC: **Receptor specificity in human, avian, and equine H2 and H3 influenza virus isolates.** *Virology* 1994, **205**:17-23.
6. Gambaryan AS, Tuzikov AB, Piskarev VE, Yamnikova SS, Lvov DK, Robertson JS, Bovin NV, Matrosovich MN: **Specification of receptor-binding phenotypes of influenza virus isolates from different hosts using synthetic sialylglycopolymers: non-egg-**

- adapted human H1 and H3 influenza A and influenza B viruses share a common high binding affinity for 6'-sialyl(N-acetyl)lactosamine). *Virology* 1997, **232**:345-350.
7. Matrosovich MN, Gambaryan AS, Teneberg S, Piskarev VE, Yamnikova SS, Lvov DK, Robertson JS, Karlsson KA: **Avian influenza A viruses differ from human viruses by recognition of sialyloligosaccharides and gangliosides and by a higher conservation of the HA receptor-binding site.** *Virology* 1997, **233**:224-234.
 8. Ito T, Couceiro JN, Kelm S, Baum LG, Krauss S, Castrucci MR, Donatelli I, Kida H, Paulson JC, Webster RG, Kawaoka Y: **Molecular basis for the generation in pigs of influenza A viruses with pandemic potential.** *J Virol* 1998, **72**:7367-7373.
 9. Couceiro JN, Paulson JC, Baum LG: **Influenza virus strains selectively recognize sialyloligosaccharides on human respiratory epithelium; the role of the host cell in selection of hemagglutinin receptor specificity.** *Virus Res* 1993, **29**:155-165.
 10. Scholtissek C: **Source for influenza pandemics.** *Eur J Epidemiol* 1994, **10**:455-458.
 11. Subbarao K, Klimov A, Katz J, Regnery H, Lim W, Hall H, Perdue M, Swayne D, Bender C, Huang J, Hemphill M, Rowe T, Shaw M, Xu XY, Fukuda Y, Cox N: **Characterization of an avian influenza A (H5N1) virus isolated from a child with a fatal respiratory illness.** *Science* 1999, **284**:393-396.
 12. Claas EC, Osterhaus AD, van Beek R, De Jong JC, Rimmelzwaan GF, Senne DA, Krauss S, Shortridge KF, Webster RG: **Human influenza A H5N1 virus related to a highly pathogenic avian influenza virus.** *Lancet* 1998, **351**:472-477.
 13. Matrosovich MN, Zhou N, Kawaoka Y, Webster R: **The surface glycoproteins of H5 influenza viruses isolated from humans, chickens, and wild aquatic birds have distinguishable properties.** *J Virol* 1999, **73**:1146-1155.
 14. Matrosovich MN, Matrosovich TY, Gray T, Roberts NA, Klenk HD: **Human and avian influenza viruses target different cell types in cultures of human airway epithelium.** *Proc Natl Acad Sci USA* 2004, **101**:4620-4624.
 15. Nicholls JM, Bourne AJ, Chen H, Guan Y, Peiris JS: **Sialic acid receptor detection in the human respiratory tract: evidence for widespread distribution of potential binding sites for human and avian influenza viruses.** *Respir Res* 2007.
 16. Matrosovich M, Krauss S, Webster R: **H9N2 influenza A viruses from poultry in Asia have human-virus-like receptor specificity.** *Virology* 2001, **281**:156-162.
 17. Saito T, Lim W, Suzuki T, Suzuki Y, Kida H, Nishimura SI, Tashiro M: **Characterization of a human H9N2 influenza virus isolated in Hong Kong.** *Vaccine* 2001, **20**:125-133.
 18. Liu J, Okazaki K, Ozaki H, Sakoda Y, Wu Q, Chen F, Kida H: **H9N2 influenza viruses prevalent in poultry in China are phylogenetically distinct from A/quail/Hong Kong/G1/97 presumed to be the donor of the internal protein genes of the H5N1 Hong Kong/97 Virus.** *Avian Pathology* 2003, **32**:551-560.
 19. Gambaryan A, Webster R, Matrosovich M: **Differences between influenza virus receptors on target cells of duck and chicken.** *Arch Virol* 2002, **147**:1197-208.
 20. Gambaryan AS, Tuzikov AB, Bovin NV, Yamnikova SS, Lvov DK, Webster RG, Matrosovich MN: **Differences between influenza virus receptors on target cells of duck and chicken and receptor specificity of the 1997 H5N1 chicken and human influenza viruses from Hong Kong.** *Avian Dis* 2003, **47**(3 Suppl):1154-1160.
 21. Gambaryan AS, Marinina VP, Solodov' TA, Bovin NV, Tuzikov AB, Pazynina GV, Yamnikova SS, Lvov DK, Klimov AI, Matrosovich MN: **Different receptor specificity of influenza viruses from ducks and chickens and its reflection in the composition of sialosides on host cells and mucins.** *Vopr Virusol* 2006, **5**:24-32.
 22. Wan H, Perez DR: **Quail carry sialic acid receptors compatible with binding of avian and human influenza viruses.** *Virology* 2006, **346**:278-286.
 23. Guo C, Takahashi N, Yagi H, Kato K, Takahashi T, Yi S, Chen Y, Ito T, Otsuki K, Kida H, Kawaoka Y, Hidari KIY, Miyamoto D, Suzuki T, Suzuki Y: **The quail and chicken intestine have sialyl-Gal sugar chains responsible for the binding of influenza A viruses to human type receptors.** *Glycobiology* 2007, **17**:713-724.
 24. Gambaryan AS, Tuzikov AB, Pazynina GV, Webster RG, Matrosovich MN, Bovin NV: **H5N1 chicken influenza viruses display a high binding affinity for the Sia2-3Galβ1-4(6-HSO₃)GlcNAc receptor.** *Virology* 2004, **326**:310-316.
 25. Gambaryan A, Tuzikov A, Pazynina G, Bovin N, Balish A, Klimov A: **Evolution of the receptor binding phenotype of influenza A (H5) viruses.** *Virology* 2006, **344**:432-438.
 26. Varki A, Cummings R, Esko J, Freeze H, Hart G, Marth J: *Essentials of Glycobiology* New York: Cold Spring Harbor Laboratory Press; 1999.
 27. Pappas C, Matsuoka Y, Swayne DE, Donis RO: **Development and Evaluation of an Influenza Virus Subtype H7N2 Vaccine Candidate for Pandemic Preparedness.** *Clin Vaccine Immunol* 2007, **14**:1425-1432.
 28. Fouchier RA, Schneeberger PM, Rozendaal FW, Broekman JM, Kemink SA, Munster V, Kuiken T, Rimmelzwaan GF, Schutten M, Van Doornum GJ, Koch G, Bosman A, Koopmans M, Osterhaus AD: **Avian influenza A virus (H7N7) associated with human conjunctivitis and a fatal case of acute respiratory distress syndrome.** *Proc Natl Acad Sci USA* 2004, **101**:1356-1361.
 29. Lee CW, Song CS, Lee YJ, Mo IP, Garcia M, Suarez DL, Kim SJ: **Sequence analysis of the hemagglutinin gene of H9N2 Korean avian influenza viruses and assessment of the pathogenic potential of isolate MS96.** *Avian Dis* 2000, **44**:527-535.
 30. Crawford PC, Dubovi EJ, Castleman WL, Stephenson I, Gibbs EPJ, Chen L, Smith C, Hill RC, Ferro P, Bright RA, Medina M, Johnson CM, Olsen CW, Cox NJ, Klimov AI, Katz JM, Donis RO: **Interspecies transmission of equine influenza virus to dogs.** *Science* 2005, **311**:1241-1242.
 31. Gambaryan A, Yamnikova S, Lvov D, Tuzikov A, Chinarev A, Pazynina G, Webster R, Matrosovich M, Bovin N: **Receptor specificity of influenza viruses from birds and mammals: new data on involvement of the inner fragments of the carbohydrate chain.** *Virology* 2005, **334**:276-283.
 32. Ha Y, Stevens DI, Skehel JJ, Wiley DC: **X-ray structure of the hemagglutinin of a potential H3 avian progenitor of the 1968 Hong Kong pandemic influenza virus.** *Virology* 2003, **309**:209-224.
 33. Russel RJ, Gamblin SJ, Haire LF, Stevens DJ, Xiao B, Ha Y, Skehel JJ: **H1 and H7 influenza haemagglutinin structures extend a structural classification of haemagglutinin subtypes.** *Virology* 2004, **325**:287-296.
 34. Lin YP, Shaw M, Gregory V, Cameron K, Lim W, Klimov A, Subbarao K, Guan Y, Krauss S, Shortridge K, Webster R, Cox N, Hay A: **Avian-to-human transmission of H9N2 subtype influenza A viruses: Relationship between H9N2 and H5N1 human isolates.** *Proc Natl Acad Sci USA* 2000, **97**:9654-9658.
 35. Vines A, Wells K, Matrosovich M, Castrucci MR, Ito T, Kawaoka Y: **The role of influenza A virus hemagglutinin residues 226 and 228 in receptor specificity and host range restriction.** *J Virol* 1998, **72**:7626-7631.
 36. Matrosovich M, Tuzikov A, Bovin N, Gambaryan A, Klimov A, Castrucci MR, Donatelli I, Kawaoka Y: **Early alterations of the receptor-binding properties of H1, H2, and H3 avian influenza virus hemagglutinins after their introduction into mammals.** *J Virol* 2000, **74**:8502-8512.
 37. Ha Y, Stevens DI, Skehel JJ, Wiley DC: **X-ray structures of H5 avian and H9 swine hemagglutinins bound to avian and human receptor analogs.** *Proc Natl Acad Sci USA* 2001, **98**:11181-11186.
 38. Yamnikova SS, Gambaryan AS, Tuzikov AB, Bovin NV, Matrosovich MN, Fedyakina IT, Grinev AA, Blinov VM, Lvov DK, Suarez DL, Swayne DE: **Differences between HA receptor-binding sites of avian influenza viruses isolated from Laridae and Anatidae.** *Avian Diseases* 2003, **47**(3 Suppl):1164-1168.
 39. Munster VJ, Baas C, Lexmond P, Waldenstrom J, Wallensten A, Fransson T, Rimmelzwaan GF, Beyer WE, Schutten M, Olsen B, Osterhaus AD, Fouchier RA: **Spatial, temporal, and species variation in prevalence of influenza A viruses in wild migratory birds.** *PLoS Pathog* 3(5):e61. 2007, May 11.
 40. Gambaryan AS, Karasin AI, Tuzikov AB, Chinarev A, Pazynina GV, Bovin NV, Matrosovich MN, Olsen CW, Klimov AI: **Receptor-binding properties of swine influenza viruses isolated and propagated in MDCK cells.** *Virus Research* 2005, **114**:15-22.
 41. Stevens J, Blixt O, Glaser L, Taubenberger JK, Palese P, Paulson JC, Wilson IA: **Glycan microarray analysis of the hemagglutinins from modern and pandemic influenza viruses reveals different receptor specificities.** *J Mol Biol* 2006, **355**:1143-1155.
 42. Matrosovich M, Matrosovich T, Uhlenhorff J, Garten W, Klenk HD: **Avian-virus-like receptor specificity of the hemagglutinin**

- impedes influenza virus replication in cultures of human airway epithelium.** *Virology* 2007, **361**:384-390.
43. Bovin NV, Korchagina EY, Zemlyanukhina TV, Byramova NE, Galanina OE, Zemlyakov AE, Ivanov AE, Zubov VP, Mochalova LV: **Synthesis of polymeric neoglycoconjugates based on N-substituted polyacrylamides.** *Glycoconj J* 1993, **10**:142-151.
 44. Gordeeva EA, Tuzikov AB, Galanina OE, Pochechueva TV, Bovin NV: **Microscale synthesis of glycoconjugate series and libraries.** *Analyt Biochem* 2000, **278**:230-232.
 45. Tuzikov AB, Gambaryan AS, Juneja LR, Bovin NV: **Conversion of complex sialooligosaccharides into polymeric conjugates and their anti-influenza virus inhibitory potency.** *J Carbohydr Chem* 2000, **19**:1191-1200.
 46. Ovchinnikova TV, Shipova EV, Sablina MA, Pazynina GA, Popova IS, Tuzikov AB, Bovin NV: **Synthesis of monosulfated saccharides in the spacers form.** *Mendeleev Commun* 2002, **12**:213-215.
 47. Pazynina GV, Sablina MA, Tuzikov AB, Chinarev AA, Bovin NV: **Synthesis of complex $\alpha(2-3)$ sialooligosaccharides, including sulfated and fucosylated ones, using Sia $\alpha(2-3)$ Gal as a building block.** *Mendeleev Commun* 2003, **13**:245-248.
 48. Gambaryan AS, Matrosovich MN: **A solid-phase enzyme-linked assay for influenza virus receptor-binding activity.** *J Virology Methods* 1992, **39**:111-123.
 49. Ng KK, Weis WI: **Structure of a selectin-like mutant of mannose-binding protein complexed with sialylated and sulfated Lewis(x) oligosaccharides.** *Biochemistry* 1997, **36**:979-988.
 50. Nicholas KB, Nicholas HB Jr, Deerfield DW: **GeneDoc: analysis and visualization of genetic variation.** *EMBNEW News* 1997, **4**:14-14.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

